


# Converging lesion and brain imaging findings uncover involvement of the cerebellum in visual perception of body motion



Pavlova et al., Perception 2001


## Neurosurgical patients (lesion-symptom mapping)






Sokolov et al., Cerebral Cortex 2010


# Healthy subjects (functional MRI)






Sokolov et al., Neurolmage 2012

Functional and effective task-specific connectivity between the left cerebellum and right superior temporal sulcus (STS), a key structure of the action observation network:






**Seed-voxel regression** 

Sokolov et al., Neurolmage 2012

# Diffusion tensor imaging (DTI) – first evidence for existence of a direct structural loop connecting the temporal cortex and cerebellum



3D reconstruction of a representative fiber tract

Sokolov et al., Cerebral Cortex 2014

#### **Conclusions**

Left lateral cerebellar structures are essential for visual processing of body motion

 First evidence in favor of direct communication and anatomical connectivity between the left cerebellum and right temporal cortex

 Potential implications for neuropsychiatric diseases such as autistic spectrum disorders, epilepsy, vertigo; and for neurorehabilitation

### In collaboration with

#### **UNIVERSITY OF TÜBINGEN MEDICAL SCHOOL**

Department of Biomedical Magnetic Resonance



**Prof. Marina Pavlova** 



Dr. Michael Erb

#### UNIVERSITY OF AACHEN MEDICAL SCHOOL

Department of Psychiatry, Psychotherapy and Psychosomatics



**Prof. Wolfgang Grodd** 

#### CENTRE HOSPITALIER UNIVERSITAIRE VAUDOIS LAUSANNE

Département des Neurosciences Cliniques



Prof. Richard Frackowiak

